

AD A 050591

EXCIMER LASER RESEARCH

J. H. Parks
AVCO EVERETT RESEARCH LABORATORY, INC.
2385 Revere Beach Parkway
Everett, MA 02149

THE VIEWS AND CONCLUSIONS CONTAINED IN THIS DOCUMENT ARE THOSE OF THE AUTHORS AND SHOULD NOT BE INTERPRETED AS NECESSARILY REPRESENTING THE OFFICIAL POLICIES, EITHER EXPRESS OR IMPLIED, OF THE DEFENSE ADVANCED RESEARCH PROJECTS AGENCY OR THE U. S. GOVERNMENT.

12

[Handwritten mark]

Semi Annual Report for Period 16 March 1977 to 15 September 1977

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.

Sponsored by
DEFENSE ADVANCED RESEARCH PROJECTS AGENCY
ARPA Order No. 1806

Monitored by
OFFICE OF NAVAL RESEARCH
DEPARTMENT OF THE NAVY
Arlington, VA 22217

DDC
RECEIVED
MAR 1 1978
F

No. 12
UDC FILE COPY

FOREWORD

DARPA Order No: 1806

Program Code No: 5E20

Name of Contractor: Avco Everett Research Laboratory, Inc.

Effective Date of Contract: 15 August 1974

Contract Expiration Date: 14 October 1977

Amount of Contract: \$762, 985

Contract No: N00014-75-C-0063

Principal Investigator and Phone No: J. H. Parks (617) 389-3000, Ext. 323

Scientific Officer: Director, Physics Program, Physical Sciences Division, Office of Naval Research, Department of the Navy, 800 North Quincy Street, Arlington, Va. 22217

Short Title of Work: Excimer Laser Research

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

048450

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)



UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

TABLE OF CONTENTS

<u>Section</u>		<u>Page</u>
	List of Illustrations	3
I.	INTRODUCTION	5
II.	E-BEAM PUMPED HgCl^* , HgBr^* LASERS	7
III.	HEAVY PARTICLE QUENCHING OF HgX^*	11
IV.	THEORETICAL EFFORT	21
V.	REFERENCES	31

ACCESSION for	
NTIS	Write Section <input checked="" type="checkbox"/>
DDC	Buff Section <input type="checkbox"/>
UNANNOUNCED	
JUSTIFICATION	
BY	DISTRIBUTION/AVAILABILITY COPIES
Dis	SPECIAL
A	

LIST OF ILLUSTRATIONS

<u>Figure</u>		<u>Page</u>
1	Schematic of Experimental Apparatus for HgX^* Quenching Kinetics	15
2	HgCl^* (B) Emission for Different Xenon Pressures	19
3	B \rightarrow X Radiative Lifetime of HgX^* as a Function of the Ionic Mixing Coefficient, β	29

I. INTRODUCTION

During this report period, research was directed towards an initial assessment of the potential for e-beam pumping of mercury monohalide lasers. Specifically, studies were undertaken to probe the molecular formation channels of HgCl in e-beam pumped mixtures of Xe/Hg/CCl_4 and Ar/Xe/Hg/CCl_4 mixtures. Consistent with this emphasis, additional experiments were begun to provide the quenching rates of the upper laser level by laser gas mixture components. Theoretical calculations of the mercury monohalide radiative lifetimes were completed which provide necessary inputs for quantitative estimates of these quenching rates as well as the molecular kinetics.

II. E-BEAM PUMPED HgCl^* , HgBr^* LASERS

During this research period the emphasis of e-beam experiments was directed towards developing a clearer understanding of the HgX^* formation kinetics and an evaluation of the laser potential. This effort included measurements of fluorescence efficiency, optimization of laser gas mixtures and output coupling, and initial measurements of HgX^* formation rates via replacement reactions involving the xenon halides. To date, pure e-beam pumping of the mercury monohalide lasers $\text{HgCl}^{*(1)}$ and $\text{HgBr}^{*(2)}$ achieved the results shown in Tables 1 and 2. Since these studies are extending beyond this research period, these results will be discussed collectively in the following semi-annual report. The relevance of these measurements to mercury monohalide e-beam pumped molecular formation kinetics will be reviewed at that time.

(1) J. H. Parks, Appl. Phys. Lett. 31, 192 (1977).

(2) J. H. Parks, Appl. Phys. Lett. 31, 297 (1977).

TABLE 1. HgCl LASER CHARACTERISTICS

• Wavelength	5576 Å
• Gas Mixture	
	Ar/Xe/Hg/CCl ₄ = 84.7%/11.1%/2.1%/1.1%
	Ar 3 Amagats
	Cell Temperature 275°C
• E-Beam Energy Deposition	4.8J
• Intrinsic Efficiency	
	(50% Coupling) 3.8%
• Peak Power	1.7 MW
• Pulsewidth	100 nsec

TABLE 2. HgBr LASER CHARACTERISTICS

• Wavelength	5018 Å
• Gas Mixture	
	Ar/Xe/Hg/HBr = 86.4%/10.8%/2.0%/.8%
	Ar 3 Amagats
	Cell Temperature 275°C
• E-Beam Energy Deposition	3 J
• Intrinsic Efficiency	
	(2.5% Coupling) .5%
• Peak Power	60 kW
• Pulsewidth	100 nsec

III. HEAVY PARTICLE QUENCHING OF HgX^*

During this past research period work was performed to design an experiment to measure the rates at which laser mixture components quench the excited state of the mercury monohalide lasers. The efficient extraction of energy from the mercury monohalide lasers will be optimized by operation at laser flux levels near the saturation flux, ϕ_s . In the presence of mixtures such as $\text{Xe}/\text{Hg}/\text{CCl}_4$, the HgCl^* molecules will be collisionally de-excited or quenched by these components. If only one of the mixture species, having a density n_Q , dominantly quenches HgCl^* at a rate k_Q , the saturation flux in the absence of bottleneaking is given by

$$\phi_s = (h\nu / \sigma_s \tau_1) (1 + \tau_1 k_Q n_Q). \quad (1)$$

Here $h\nu$ is the laser photon energy, σ_s the stimulated emission cross section, and τ_1 the HgCl^* radiative lifetime. In a real case the quenching by each mixture species will have to be taken into account, and this will probably include both two and three body quenching processes. Thus, a measurement of the relevant HgCl^* quenching kinetics is necessary for the determination of the saturation flux and efficient laser operation. A method of determining the relevant quenching processes for the mercury monohalide system, relies on optical pumping to produce HgX^* in the presence of the quenching species

In choosing the best mix for a particular laser system various components are added to perform different functions. In a discharge,

for example, components are chosen which will produce a desired electron temperature, stabilize the discharge, have large cross sections for forming the desired upper lasing level, have small optical absorption at the lasing frequency, etc. All these components, however, can, in addition to their desired effect, cause the unintentional quenching of the upper laser level. Quenching can be caused by both the heavy particles and the electrons. It is the purpose of this research to measure the effect of the heavy particles on the upper HgX^* laser level. In the following experimental scheme HgX^* will be produced by optical pumping of the mercury dihalide salt (HgX_2) thus eliminating any ambiguity in the interpretation of the results due to the presence of electrons. This should allow for straightforward interpretation of the data.

The work of Wieland⁽³⁾ indicates that one can obtain emission from the mercury monohalide excited states (HgX^*) by pumping the stable mercury dihalide salts with ultraviolet light. Wieland has shown that the pump wavelength ranges given in Table 3 optimally produce HgX^* . The range is determined by those wavelengths which restrict the photodissociation channel of HgX_2 to that producing the mercury monohalide state HgX^* ($B^2\Sigma_{1/2}^+$). For the mercury chloride system, Wieland's work indicates that if HgCl_2 is optically pumped by 1700 Å radiation, one observes HgCl^* ($B \rightarrow X$) emission. The Xe_2^* excimer is a well known bright fluoressor with peak emission at 1720 Å. Thus, if one were to use a xenon lamp to pump HgCl_2 , one would produce adequate quantities of HgCl^* for these quenching measurements.


(3) K. Wieland, Zeits. f. Physik 77, 157 (1932).

TABLE 3. HgX_2 ABSORPTION WAVELENGTHS
FOR $\text{HgX}^*(B^2\Sigma^+_{1/2})$ PRODUCTION

	<u>λ min (\AA)</u>	<u>λ max (\AA)</u>
HgCl	1600	1810
HgBr	1700	1950
HgI	1937	2254

In order to pursue this course of investigation we have designed and built a Π shaped quartz gas discharge tube which is filled with ~ 200 torr xenon gas to produce Xe_2^* emission. This design is based on the work of Huffman et al. (4) A schematic diagram of the discharge tube and associated experimental equipment is shown in Figure 1. The Xe_2^* emission is coupled into the reaction cell which is heated in a double oven configuration. The HgX_2 salt is heated at one temperature (T_1) in a separate side arm which serves to control the HgX_2 vapor pressure. For HgCl^* measurements, a vapor pressure of $\text{HgCl}_2 \sim 50$ microns is adequate ($T_1 \approx 90^\circ\text{C}$). The main body of the reaction cell is kept at a temperature (T_2) which is higher than T_1 and ensures that the windows remain clear. In this configuration, the vapor pressure and HgCl_2 density in the cell is accurately known; for example if $T_2 \approx 200^\circ\text{C}$, we have $n(\text{HgCl}_2) \approx 10^{15} \text{ cm}^{-3}$. A 1/2 m/JA monochromator and PM1 is used to record the HgX^* spectra through the side of the cell and since the quartz will not pass the Xe_2^* radiation, only the HgX^* emission will be detected. The lamp is pulsed at 300-400 Hz and a boxcar integrator is used to improve signal to noise. The visible Xe discharge fluorescence is monitored (PM2) at one end of the lamp to check the general behavior of the discharge. The observed reproducibility of the HgCl^* emission assures that the Xe_2^* fluorescence is itself quite reproducible. When taking quenching rate data the HgCl^* fluorescence signal is directly monitored by PM3. A filter for Xe_2^* pump radiation is positioned between the discharge lamp and the reaction cell in order to reduce visible Xe background emission. An additional cell can also be placed between the

(4) R. E. Huffman et al, Appl. Optics 2, 617 (1963).







H1371

Figure 1 Schematic of Experimental Apparatus for HgX^* Quenching Kinetics

lamp and cell to measure the possible ultraviolet absorption by the quenching species. In this case the visible HgX^* emission indicates the ultraviolet transmission thus avoiding the need for ultraviolet measurements.

An outline of the HgCl^* kinetics and experimental analysis for this scheme follow from the reactions:



where I is the intensity of the Xe_2^* radiation exciting the HgCl_2 . These lead to the rate equation

$$\frac{d[\text{HgCl}^*]}{dt} = k_I I [\text{HgCl}_2] - k_Q [\text{HgCl}^*] [\text{Q}] - \frac{[\text{HgCl}^*]}{\tau} - k' [\text{HgCl}^*] [\text{HgCl}_2] \quad (2)$$

which has a solution under steady state conditions (i. e., $\frac{d[\text{HgCl}^*]}{dt} = 0$) is given by

$$[\text{HgCl}^*] = \frac{k_I I [\text{HgCl}_2]}{k_Q [\text{Q}] + 1/\tau + k' [\text{HgCl}_2]} \quad (3)$$

The detected HgCl^* emission signal is proportional to

$$S_Q = \frac{[\text{HgCl}^*]}{\tau} = \frac{k_I I [\text{HgCl}_2]}{1 + k_Q [Q] \tau + k' [\text{HgCl}_2] \tau} \quad (4)$$

Similarly if $[Q] = 0$, we have

$$S_0 = \frac{k_I I [\text{HgCl}_2]}{1 + k' [\text{HgCl}_2] \tau} \quad (5)$$

Finally, define the ratio R by

$$R = \frac{S_0}{S_Q} = 1 + \frac{k_Q Q}{1/\tau + k' [\text{HgCl}_2]} \quad (6)$$

For expected conditions of the experiment

$$\begin{aligned} k_Q &\leq 2 \times 10^{-10} \text{ cm}^3/\text{sec} \text{ (slightly less than gas kinetic)} \\ \tau (\text{HgCl}^*) &\simeq 3 \times 10^{-8} \text{ sec (Ref. 5)} \\ [\text{HgCl}_2] &\gtrsim 10^{15} \text{ cm}^{-3} \end{aligned} \quad (7)$$

we observe that $1/\tau \gg k' [\text{HgCl}_2]$, i. e. the quenching by the HgCl_2 background is negligible and R is directly proportional to the quenching rate K_Q

$$R = 1 + k_Q [Q] \tau$$

(5) N. Djeu and C. Mazza, Chem. Phys. Lett. 46, 172 (1977).

Thus a Stern-Volmer plot of R vs $[Q]$ yields the slope $k_Q \tau$. Plots of this type will be made for the different mercury monohalides as a function of relevant quenching gases.

During this past research period the apparatus for this experiment was designed and assembled. A preliminary experiment was performed to gain a better understanding of the utility of this technique. In this experiment the HgCl^* (B - X) spectrum was recorded as a function of xenon pressure. The HgCl^* spectrum for a xenon pressure of 200 torr is compared with a spectrum in the absence of xenon in Figure 2. Planimeter measurements show comparable areas (total energy emitted) for these two cases which indicates negligible HgCl^* quenching at this pressure. However, the drastic change in the spectra arises from the rapid relaxation of the HgCl^* vibrational levels by Xe. When $[\text{Xe}] = 0$, the HgCl^* spectrum extends to roughly 3750 \AA which is identified with transitions from $B^2\Sigma_{1/2}^+$ levels $v' \sim 20 - 25$ to $X^2\Sigma_{1/2}^+$ levels $v'' \sim 0 - 5$. This HgCl^* photon energy of $\sim 3.3 \text{ eV}$ corresponds roughly to the Xe_2^* photon energy (6.8 eV) less the $\text{HgCl}-\text{Cl}$ bond energy of 3.5 eV . This indicates that high vibrational levels of the B state are produced in the photodissociation of HgCl_2 by Xe_2^* radiation. For $[\text{Xe}] = 200 \text{ torr}$, the excited state vibrational manifold is collisionally relaxed and primarily the strong $v' = 0 \rightarrow v'' = 22 - 15$ radiative transitions are observed. This relaxation can be quite rapid since the vibrational energy spacing is $\sim 190 \text{ cm}^{-1} \leq kT$. Comparable results are also observed in HgCl^* laser spectra⁽¹⁾ resulting from mixtures containing a xenon density of $\sim 1 \text{ amagat}$.

Preliminary data indicates a two-body xenon quenching rate of $k_{\text{Xe}} \sim 3 \times 10^{-13} \text{ cm}^3 \text{ sec}^{-1}$. A most important result indicated by this




Figure 2 HgCl^* (B) Emission for Different Xenon Pressures

measurement is that attempts to measure the quenching by molecular species such as CCl_4 at low pressures will require an admixture with about 200 torr of xenon to insure that the quenching rate measurement primarily represents the quantity of interest: de-excitation from the $v' = 0$ HgCl^* laser level.

This xenon density does not introduce significant quenching since $[\text{Xe}] k_{\text{Xe}} \tau \approx .06 \ll 1$ at $[\text{Xe}] = 200$ torr which makes such a procedure reliable.

IV. THEORETICAL EFFORT

This section will discuss the application of a charge transfer model to calculate the oscillator strengths and radiative lifetimes for the mercury-monohalide laser transitions $B^2\Sigma_{1/2}^+ - X^2\Sigma_{1/2}^+$. To our knowledge, no previous calculations of the radiative lifetimes of these systems have been attempted. Dunning and Hay⁽⁶⁾ have performed ab initio calculations for KrF, in which they computed both the potential energy curves and oscillator strengths. Their calculation involved an extensive configuration-interaction treatment, utilizing between 2000-3000 configurations. Although such calculations are valuable, they are enormously time consuming and must be applied individually to each state of each rare gas-halide pair. This approach is clearly not in a form from which general conclusions can be drawn and applied to the whole class of rare gas-halide systems. It would therefore be desirable to have available a simpler model, from which one can make fairly reliable predictions for a wide range of systems. The simple charge transfer theory developed originally by Mulliken⁽⁷⁾ and used by Zare and Herschbach⁽⁸⁾ for the alkali halides, appears to be a scheme capable of being generalized to a wide variety of systems for which the "ionic-bonding" model holds, including the alkali halides, rare gas-halides, and mercury-monohalides. The alkali-halide analogy has been extremely

(6) P. Hay and T. Dunning Jr., J. Chem. Phys. 66, 1306 (1977).

(7) R. S. Mulliken, J. Chem. Phys. 7, (1939) 20.

(8) R. N. Zare and D. R. Herschbach, J. Mol. Spectry. 15 (1965) 462.

successful⁽⁹⁾ in predicting the properties of the rare gas-halides, and the use of the charge transfer model to calculate the transition strengths can be thought of as simply an extension of the analogy.

The basic idea behind the charge transfer theory is that the electron initially localized around the halogen center, in the upper M^+X^- state, "jumps" or is transferred to the mercury positive ion, filling the p-state vacancy, and thus ending up in the MX atomic ground state. This is essentially a "valence bond" picture, in which the electron is transferred from one atomic center to the other. For example, in the mercury chloride $B \rightarrow X$ transition an electron which is initially localized on the Cl atom, forming Cl^- , is transferred to the Hg^+ ion resulting in the ground state configuration of mercury chloride.

In general, the mercury-halides are closely analogous in structure to the rare gas-halides. The upper laser level is "ionic", while the ground state is primarily "atomic". There are, however, some significant differences between the two systems. The upper level of the mercury-halides correlates with the $Hg^+(^2S_{1/2}) + X^-(^1S_0)$ atomic states so that only a single $^2\Sigma_{1/2}$ ionic state is formed. The ground state correlates with the $Hg(^1S_0) + X(^2P_{3/2}, ^1/2)$ atomic limit, so that we obtain $^2\Pi_{1/2}$, and $^2\Sigma_{1/2}$ molecular states, as in the case of the rare gas-halides. A more important difference is that the ground states of the mercury-halides are essentially bound states with the binding energy ranging from⁽¹⁰⁾ .36 eV for HgI to 1.8 eV for HgF. With the exception of XeF, which is slightly bound, all of the rare gas-halide ground states are repulsive.

(9) J. J. Ewing, C. A. Brau, Phys. Rev. A12, 129 (1975).

(10) G. Herzberg, Spectra of Diatomic Molecules, 2nd Ed. (Van Nostrand, Princeton, 1950).

Although the upper and lower states are predominantly ionic and covalent, respectively, it proves to be essential to allow for mixing between these states in order to obtain reasonable results from the theory. The mixed X and B state wavefunctions can be expressed as

$$\begin{aligned}\Psi_B &= N_B (\Psi^i + \alpha \Psi^c), \\ \Psi_X &= N_X (\Psi^c - \beta \Psi^i),\end{aligned}\tag{8}$$

with $\alpha = (\beta - S)/(1 - \beta S)$, and where Ψ^i and Ψ^c are the purely ionic and covalent wavefunctions, N_B and N_X are normalization constants, S is the overlap integral between Ψ^i and Ψ^c , and β is the mixing coefficient. Except for very large values of the internuclear distance, S is non-zero so that S as well as β must appear in the expression for α in order to make the initial and final wavefunctions orthogonal. A major difficulty in the theory is the specification of the mixing parameter β . Given the lack of data on the mercury-monohalides, we have used the procedure outlined by Coulson⁽¹¹⁾ to estimate the degree of ionic character of the ground state from the electronegativities of the mercury and halogen atoms. The values for the percent ionic character [i. e. $100 \beta^2/(1 + \beta^2)$] obtained in this way are given in Table 4. The electronegativities can also be used to determine the co-called covalent-ionic resonance energy, Δ , which is simply the ionic contribution to the binding energy. The values of Δ (in eV) turn out to be 1.2, 0.81, and 0.36 for HgCl, HgBr, and HgI, respectively, as compared to the experimental dissociation energies (in eV) of 1.0, 0.7, and 0.36. It

(11) C. A. Coulson, *Valence*, 2nd Ed. (Oxford Univ. Press, London, 1961) pp. 139-141.

TABLE 4. PARAMETERS FOR THE B → X TRANSITION IN THE Hg-HALIDES

	α	β	S	$X^2\Sigma$ Percent Ionic Character	$B^2\Sigma$ Internuclear Equilibrium Distance(\AA)	$\lambda(\text{\AA})$	$\mu(\text{Debye})$	τ (nsec)		
								Theory	Expt.	Scaled
HgCl	.64	.53	.87	22	3.15	5576	-6.60	20	--	29
HgBr	.59	.45	.82	17	3.24	5018	-6.32	16	23*	23
HgI	.51	.35	.73	11	3.50	4412	-5.92	12	--	17

* Reference 5

may thus be inferred that the electronegativity method tends to overestimate the mixing parameter.

In order to complete the definition of the wavefunctions in Eq. 8, it remains to determine the one-electron orbitals that make up Ψ^c and Ψ^i . In keeping with the valence bond picture we have adopted, Ψ^c and Ψ^i are simply taken to be products of unperturbed atomic (ionic) wavefunctions. The analytic Hartree-Fock functions of Clementi and Roetti⁽¹²⁾ were used for the atomic and ionic halogen wavefunctions. Unfortunately, no such functions exist for neutral or singly-ionized mercury. A further complication arises from the fact that relativistic effects are non-negligible for the mercury valence orbital. Since the valence electrons are in an *s*-state, the most important relativistic effect⁽¹³⁾ is a contraction of the radial wavefunction toward the nucleus, due to the absence of a repulsive angular momentum barrier. We have therefore calculated a non-relativistic Hartree-Fock-Slater (HFS) wavefunction, utilizing the Herman-Skillman computer program⁽¹⁴⁾, and scaled the function according to the formula

$$P'(r) = \gamma^{1/2} P_{\text{non-rel.}}(\gamma r)$$

so as to reproduce the expectation values $\langle r^n \rangle$ tabulated by Lu et al,⁽¹⁵⁾ from their relativistic HFS program. The calculation was made for $n = -1, 1, 2$ and in all cases $\gamma \approx 1.16$ for Hg; the same value was used for Hg^+ . The final function, $P'(r)$, were then represented by a sum of

(12) E. Clementi and C. Roetti, *At. Nucl. Data Tables* 14 (1974) 177.

(13) V. M. Burke and I. P. Grant, *Proc. Phys. Soc. (London)* 90 (1967) 297.

(14) F. Herman and S. Skillman, *Atomic Structure Calculations* (Prentice-Hall, Englewood Cliffs, 1963).

(15) C. C. Lu, T. A. Carlson, F. B. Malik, T. C. Tucker and C. W. Nestor Jr., *At. Data* 3 (1971) 1.

Slater-type orbitals (STO's) using a nonlinear least squares fitting procedure. The basis set for the 6s functions of Hg and Hg^+ consisted of eight STO's.

With the wavefunctions given in Eq. (8), the transition dipole moment can be reduced to an expression involving only one-electron terms. Consider that the ionic and covalent molecular wavefunctions Ψ^i and Ψ^c are expressed in terms of products of atomic wavefunctions Φ for Hg, Hg^+ , X and X^- ($X = \text{Cl}, \text{Br}, \text{I}$) by

$$\begin{aligned}\Psi^i &= (\Phi_{\text{Hg}^+}) (\Phi_{X^-}) \\ \Psi^c &= (\Phi_{\text{Hg}}) (\Phi_X).\end{aligned}\tag{9}$$

These atomic wavefunctions Φ are represented by Hartree-Fock functions composed of products of one-electron wavefunctions, ϕ . Then, the transition dipole moment μ can be shown to be given by

$$\begin{aligned}\mu &= N_B N_X [(\phi_{\text{Hg}} | r | \phi_{X^-}) (1 - \alpha\beta)\chi \\ &\quad + \alpha \langle \phi_{\text{Hg}} | r | \phi_{\text{Hg}} \rangle - \beta \langle \phi_{X^-} | r | \phi_{X^-} \rangle],\end{aligned}\tag{10}$$

where ϕ_{X^-} is the initial wavefunction of the active electron centered on the negative halogen ion (X^-), ϕ_{Hg} is the final state one-electron wavefunction, r is the position vector, and χ is an overlap integral involving all of the electron orbitals not involved in the transition. All two-center integrals were evaluated from an expansion developed by Sharma⁽¹⁶⁾ for

(16) R. R. Sharma, Phys. Rev. A13 (1976) 517.


the case of one-electron wavefunctions expressed as a sum of STO's. The calculation of the overlap function $S \equiv \langle \Psi^i | \Psi^c \rangle$ and the parameter α for various mercury monohalides are given in Table 4. All these calculations depend on a choice of molecular internuclear separation. An experimental value for the equilibrium internuclear distance exists⁽¹⁷⁾ for the ionic state of HgCl, but not for the other halides. In order to estimate these distances, the relative sizes of the alkali halides were used as a guide. It was found that the internuclear separation of the alkali bromides are $\approx 3\%$ larger than those of the alkali chlorides while the iodides are $\approx 12\%$ larger. The internuclear distances of the Hg-halides were scaled accordingly, and are given in Table 4.

The results of the calculations for the $B \rightarrow X$ transitions are summarized in Table 4. The radiative lifetime of the transition is given, in terms of the quantity μ , by $\tau = 3h\lambda^3/64\pi^4\mu^2$ where λ is the transition wavelength. The model predicts a decrease in lifetime with increasing halogen size due to the λ^3 -dependence. There is a competing trend caused by a decrease in the dipole matrix element, but this effect is much weaker. In Table 4 we give the experimental result of Djeu and Mazza.⁽⁵⁾ The theoretical value is $\approx 30\%$ lower than the measured lifetime, which is quite satisfactory considering the simplicity of the model. Further calculations were carried out to test the sensitivity of the results to the various parameters. When the internuclear distance was varied by 5%, the lifetime changed by less than 5% in all cases. As discussed earlier, the mixing coefficients are the parameters which contain the greatest uncertainty. To give some idea of the sensitivity of our calculation to the percent ionic

(17) K. Wieland, *Helv. Phys. Acta* 14 (1941) 420.

character of the ground state, the radiative lifetime of each mercury monohalide is plotted as a function of the mixing coefficient in Figure 3. The value of the mixing coefficient inferred from the electronegativity is indicated for each molecule in Figure 3 by the cross. From our earlier arguments, we expect these mixing coefficients in Table 4 to be over-estimates, so that from Figure 3 the calculated lifetimes are expected to be somewhat too small. This is consistent with the comparison between theory and experiment for HgBr. The sensitivity of the radiative lifetime to a variation of the mixing coefficient β can be estimated from Figure 3. It is easily shown that the variation of a ratio of lifetimes is less sensitive to β and thus a scaling of the calculated lifetimes to the HgBr experimental value is a useful estimate. These scaled lifetimes are also shown in Table 4. Finally, an estimate of the importance of effects such as HgX^* polarization and the broadband emission spectrum were considered and are included in Ref. 18.

(18) C. Duzy, H. A. Hyman, Chem. Phys. Lett. 52, 345 (1977).



H3733

Figure 3 $B \rightarrow X$ Radiative Lifetime of HgX^* as a Function of the Ionic Mixing Coefficient, β

V. REFERENCES

1. J. H. Parks, Appl. Phys. Lett. 31, 192 (1977).
2. J. H. Parks, Appl. Phys. Lett. 31, 297 (1977).
3. K. Wieland, Zeits. f. Physik 77, 157 (1932).
4. R. E. Huffman et al, Appl. Optics 2, 617 (1963).
5. N. Djeu and C. Mazza, Chem. Phys. Lett. 46, 172 (1977).
6. P. Hay and T. Dunning Jr., J. Chem. Phys. 66, 1306 (1977).
7. R. S. Mulliken, J. Chem. Phys. 7 (1939) 20.
8. R. N. Zare and D. R. Herschbach, J. Mol. Spectry. 15 (1965) 462.
9. J. J. Ewing, C. A. Brau, Phys. Rev. A12, 129 (1975).
10. G. Herzber, Spectra of Diatomic Molecules, 2nd Ed. (Van Nostrand, Princeton, 1950).
11. C. A. Coulson, Valence, 2nd Ed. (Oxford Univ. Press, London, 1961) pp. 139-141.
12. E. Clementi and C. Roetti, At. Nucl. Data Tables 14 (1974) 177.
13. V. M. Burke and I. P. Grant, Proc. Phys. Soc. (London) 90 (1967) 297.
14. F. Herman and S. Skillman, Atomic Structure Calculations (Prentice-Hall, Englewood Cliffs, 1963).
15. C. C. Lu, T. A. Carlson, F. B. Malik, T. C. Tucker and C. W. Nestor Jr., At. Data 3 (1971) 1.
16. R. R. Sharma, Phys. Rev. A13 (1976) 517.
17. K. Wieland, Helv. Phys. Acta 14 (1941) 420.
18. C. Duzy, H. A. Hyman, Chem. Phys. Lett. 52, 345 (1977).

DISTRIBUTION LIST

Office of Naval Research, Department of the Navy, Arlington, VA 22217 - Attn: Physics Program (3 copies)

Naval Research Laboratory, Department of the Navy, Washington, D.C. 20375 - Attn: Technical Library (1 copy)

Office of the Director of Defense, Research and Engineering, Information Office Library Branch, The Pentagon, Washington, D.C. 20301 (1 copy)

U.S. Army Research Office, Box CM, Duke Station, Durham, N.C. 27706 (1 copy)

Defense Documentation Center, Cameron Station, Alexandria, VA 22314 (12 copies)

Defender Information Analysis Center, Battelle Memorial Institute, 505 King Avenue, Columbus, OH 43201 (1 copy)

Commanding Officer, Office of Naval Research Branch Office, 536 South Clark Street, Chicago, IL 60615 (1 copy)

New York Area Office, Office of Naval Research, 715 Broadway (5th Floor), New York, NY 10003 - Attn: Dr. Irving Rowe (1 copy)

San Francisco Area Office, Office of Naval Research, 760 Market Street, Room 447, San Francisco, CA 94102 (1 copy)

Air Force Office of Scientific Research, Department of the Air Force, Washington, D.C. 22209 (1 copy)

Office of Naval Research Branch Office, 1030 East Green Street, Pasadena, CA 91106 - Attn: Dr. Robert Behringer (1 copy)

Code 102 1P (ONRL), Office of Naval Research, 800 N. Quincy Street, Arlington, VA 22217 (6 copies)

Defense Advanced Research Projects Agency, 1400 Wilson Blvd., Arlington, VA 22209 - Attn: Strategic Technology Office (1 copy)

Office Director of Defense, Research & Engineering, The Pentagon, Washington, D.C. 20301 - Attn: Assistant Director (Space and Advanced Systems) (1 copy)

Office of the Assistant Secretary of Defense, System Analysis (Strategic Programs), Washington, D.C. 20301 - Attn: Mr. Gerald R. McNichols (1 copy)

U.S. Arms Control and Disarmament Agency, Dept. of State Bldg., Rm. 4931, Washington, D.C. 20451 - Attn: Dr. Charles Henkin (1 copy)

Energy Research Development Agency, Division of Military Applications, Washington, D.C. 20545 (1 copy)

National Aeronautics and Space Administration, Lewis Research Center, Cleveland, OH 44135 - Attn: Dr. John W. Dunning, Jr. (1 copy)
(Aerospace Res. Engineer)

National Aeronautics & Space Administration, Code RR, FOB 10B, 600 Independence Ave., SW, Washington, D.C. 20546 (1 copy)

National Aeronautics and Space Administration, Ames Research Center, Moffett Field, CA 94035 - Attn: Dr. Kenneth W. Billman (1 copy)

Department of the Army, Office of the Chief of RD&A, Washington, D.C. 20310 - Attn: DARD-DD (1 copy)
DAMA-WSM-T (1 copy)

Department of the Army, Office of the Deputy Chief of Staff for Operations & Plans, Washington, D.C. 20310 - Attn: DAMO-RQD - (1 copy)

Ballistic Missile Defense Program Office (BMDPO), The Commonwealth Building, 1300 Wilson Blvd., Arlington, VA 22209 - Attn: Mr. Albert J. Bast, Jr. (1 copy)

U.S. Army Missile Command, Research & Development Division, Redstone Arsenal, AL 35809 - Attn: Army High Energy Laser Programs (2 copies)

Commander, Rock Island Arsenal, Rock Island, IL 61201, Attn: SARRI-LR, Mr. J. W. McGarvey (1 copy)

Commanding Officer, U.S. Army Mobility Equipment R&D Center, Ft. Belvoir, VA 22060 - Attn: SMEFB-MW (1 copy)

Commander, U.S. Army Armament Command, Rock Island, IL 61201 - Attn: AMSAR-RDT (1 copy)

Director, Ballistic Missile Defense Advanced Technology Center, P.O. Box 1500, Huntsville AL 35807 - Attn: ATC-O (1 copy)
ACT-T (1 copy)

Commander, U.S. Army Material Command, Alexandria, VA 22304 - Attn: Mr. Paul Chernoff (AMCRD-T) (1 copy)

Commanding General, U.S. Army Munitions Command, Dover, NH 17801 - Attn: Mr. Gilbert F. Chesnov (AMSMU-R) (1 copy)

Director, U.S. Army Ballistics Res. Lab, Aberdeen Proving Ground, MD 21005 - Attn: Dr. Robert Eichenberger (1 copy)

Commandant, U.S. Army, Air Defense School, Ft. Bliss, TX 79916 - Attn: Air Defense Agency (1 copy)
ATSA-CTD-MS (1 copy)

Commanding General, U.S. Army Combat Dev. Command, Ft. Belvoir, VA 22060 - Attn: Director of Material, Missile Div. (1 copy)

Commander, U.S. Army Training & Doctrine Command, Ft. Monroe, VA 23651 - Attn: ATCD-CF (1 copy)

Commander, U.S. Army Frankford Arsenal, Philadelphia, PA 19137 - Attn: Mr. M. Elnick SARFA-FCD Bldg. 201-3 (1 copy)

Commander, U.S. Army Electronics Command, Ft. Monmouth, NJ 07703 - Attn: AMSEL-CT-L, Dr. R.G. Buser (1 copy)

Commander, U.S. Army Combined Arms Combat Developments Activity, Ft. Leavenworth, KS 66027 (1 copy)

National Security Agency, Ft. Geo. G. Meade, MD 20755 - Attn: R.C. Foss A763 (1 copy)

Deputy Commandant for Combat & Training Developments, U.S. Army Ordnance Center and School, Aberdeen Proving Ground, MD 21005
Attn: ATSL-CTD-MS-R (1 copy)

Commanding Officer, USACDC CBR Agency, Ft. McClellan, AL 36201 - Attn: CDCCBR-MR (Mr. F.D. Poer) (1 copy)

BEST AVAILABLE COPY

BEST AVAILABLE COPY

DISTRIBUTION LIST (Continued)

Department of the Navy, Office of the Chief of Naval Operations, The Pentagon 5C739, Washington, D.C. 20350 - Attn: (OP 982F3) (1 copy)

Office of Naval Research Branch Office, 495 Summer Street, Boston, MA 02210 - Attn: Dr. Fred Quelle (1 copy)

Department of the Navy, Deputy Chief of Navy Material (Dev.), Washington, D.C. 20360 - Attn: Mr. R. Gaylord (MAT 032B) (1 copy)

Naval Missile Center, Point Mugu, CA 93042 - Attn: Gary Gibbs (Code 5352) (1 copy)

Naval Research Laboratory, Washington, D.C. 20375 - Attn: (Code 5503-EOTPO) (1 copy)
 Dr. P. Livingston - Code 5560 (1 copy)
 Dr. A. I. Schindler - Code 6000 (1 copy)
 Dr. H. Shenker - Code 5504 (1 copy)
 Mr. D. J. McLaughlin - Code 5560 (1 copy)
 Dr. John L. Walsh - Code 5503 (1 copy)

High Energy Laser Project Office, Department of the Navy, Naval Sea Systems Command, Washington, D.C. 20360 - Attn: Capt. A. Skolnick, USN (PM 22) (1 copy)

Superintendent, Naval Postgraduate School, Monterey, CA 93940 - Attn: Library (Code 2124) (1 copy)

Navy Radiation Technology, Air Force Weapons Lab (NLO), Kirtland AFB, NM 87117 (1 copy)

Naval Surface Weapons Center, White Oak, Silver Spring, MD 20910 - Attn: Dr. Leon H. Schindel (Code 310) (1 copy)
 Dr. E. Leroy Harris (Code 313) (1 copy)
 Mr. K. Enkenhaus (Code 034) (1 copy)
 Mr. J. Wise (Code 047) (1 copy)
 Technical Library (1 copy)

U.S. Naval Weapons Center, China Lake, CA 93555 - Attn: Technical Library (1 copy)

HQ USAF (AF/RDPS), The Pentagon, Washington, D.C. 20330 - Attn: Lt. Col. A. J. Chiota (1 copy)

HQ AFSC/XRLW, Andrews AFB, Washington, D.C. 20331 - Attn: Maj. J. M. Walton (1 copy)

HQ AFSC (DLCAW), Andrews AFB, Washington, D.C. 20331 - Attn: Maj. H. Axelrod (1 copy)

Air Force Weapons Laboratory, Kirtland AFB, NM 87117 - Attn: LR (1 copy)
 AL (1 copy)

HQ SAMSO (XRTD), P.O. Box 92960, Worldway Postal Center, Los Angeles, CA 90009 - Attn: Lt. Dorian DeMaio (XRTD) (1 copy)

AF Avionics Lab (TEO), Wright Patterson AFB, OH 45433 - Attn: Mr. K. Hutchinson (1 copy)

Dept. of the Air Force, Air Force Materials Lab. (AFSC), Wright Patterson AFB, OH 45433 - Attn: Maj. Paul Elder (LPS) (1 copy)
 Laser Window Group

HQ Aeronautical Systems Div., Wright Patterson AFB, OH 45433 - Attn: XRF - Mr. Clifford Fawcett (1 copy)

Rome Air Development Command, Griffiss AFB, Rome, NY 13440 - Attn: Mr. R. Urtz (OCSE) (1 copy)

HQ Electronics Systems Div. (ESL), L. G. Hanscom Field, Bedford, MA 01730 - Attn: Mr. Alfred E. Anderson (XRT) (1 copy)
 Technical Library (1 copy)

Air Force Rocket Propulsion Lab., Edwards AFB, CA 93523 - Attn: B. R. Bornhorst, (LKCG) (1 copy)

Air Force Aero Propulsion Lab., Wright Patterson AFB, OH 45433 - Attn: Col. Walter Moe (CC) (1 copy)

Dept. of the Air Force, Foreign Technology Division, Wright Patterson AFB, OH 45433 - Attn: PDTN (1 copy)

Commandant of the Marine Corps, Scientific Advisor (Code RD-1), Washington, D.C. 20380 (1 copy)

Aerospace Research Labs., (AP), Wright Patterson AFB, OH 45433 - Attn: Lt. Col. Max Duggins (1 copy)

Defense Intelligence Agency, Washington, D.C. 20301 - Attn: Mr. Seymour Berler (DT1B) (1 copy)

Central Intelligence Agency, Washington, D.C. 20505 - Attn: Mr. Julian C. Nall (1 copy)

Analytic Services, Inc., 5613 Leesburg Pike, Falls Church, VA 22041 - Attn: Dr. John Davis (1 copy)

Aerospace Corp., P.O. Box 92957, Los Angeles, CA 90009 - Attn: Dr. G. P. Millburn (1 copy)

Airresearch Manuf. Co., 9851-9951 Sepulveda Blvd., Los Angeles, CA 90009 - Attn: Mr. A. Colin Stancliffe (1 copy)

Atlantic Research Corp., Shirley Highway at Edsall Road, Alexandria, VA 22314 - Attn: Mr. Robert Naismith (1 copy)

Avco Everett Research Lab., 2385 Revere Beach Parkway, Everett, MA 02149 - Attn: Dr. George Sutton (1 copy)
 Dr. Jack Daugherty (1 copy)

Battelle Columbus Laboratories, 505 King Avenue, Columbus, OH 43201 - Attn: Mr. Fred Tietzel (STPIAC) (1 copy)

Bell Aerospace Co., Buffalo, NY 14240 - Attn: Dr. Wayne C. Solomon (1 copy)

Boeing Company, P.O. Box 3999, Seattle, WA 98124 - Attn: Mr. M. I. Gamble (2-, 460, MS 8C-88) (1 copy)

Electro-Optical Systems, 300 N. Halstead, Pasadena, CA 91107 - Attn: Dr. Andrew Jensen (1 copy)

ESL, Inc., 495 Java Drive, Sunnyvale, CA 94086 - Attn: Arthur Einhorn (1 copy)

BEST AVAILABLE COPY

DISTRIBUTION LIST (Continued)

General Electric Co., Space Division, P.O. Box 8555, Philadelphia, PA 19101 - Attn: Dr. R. R. Sigismonti (1 copy)
General Electric Co., 100 Plastics Avenue, Pittsfield, MA 01201 - Attn: Mr. D. G. Harrington (Rm. 1044) (1 copy)
General Research Corp., P.O. Box 3587, Santa Barbara, CA 93105 - Attn: Dr. R. Holbrook (1 copy)
General Research Corp., 1501 Wilson Blvd., Suite 700, Arlington, VA 22209 - Attn: Dr. Giles F. Crimi (1 copy)
Hercules, Inc., Industrial System Dept., Wilmington, DE 19899 - Attn: Dr. R. S. Voris (1 copy)
Hercules, Inc., P.O. Box 210, Cumberland, MD 21502 - Attn: Dr. Ralph R. Preckel (1 copy)
Hughes Research Labs., 3011 Malibu Canyon Road, Malibu, CA 90265 - Attn: Dr. D. Forster (1 copy)
Hughes Aircraft Co., Aerospace Group - Systems Division, Canoga Park, CA 91304 - Attn: Dr. Jack A. Alcalay (1 copy)
Hughes Aircraft Co., Centinela and Teale Streets, Bldg. 6, MS E-125, Culver City, CA 90230 - Attn: Dr. William Yates (1 copy)
Institute for Defense Analyses, 400 Army-Navy Drive, Arlington, VA 22202 - Attn: Dr. Alvin Schnitzler (1 copy)
Johns Hopkins University, Applied Physics Lab., 8621 Georgia Avenue, Silver Spring, MD 20910 - Attn: Dr. Albert M. Stone (1 copy)
Lawrence Livermore Laboratory, P.O. Box 808, Livermore, CA 94550 - Attn: Dr. R. E. Kidder (1 copy)
Dr. E. Teller (1 copy)
Dr. Joe Fleck (1 copy)
Los Alamos Scientific Laboratory, P.O. Box 1663, Los Alamos, NM 87544 - Attn: Dr. Keith Boyer (1 copy)
Lulejian and Associates, Inc., Del Amo Financial Center, 21515 Hawthorne Blvd. - Suite 500, Torrance, CA 90503 (1 copy)
Lockheed Palo Alto Res. Lab., 3251 Hanover St., Palo Alto, CA 94303 - Attn: L. R. Lunsford, Orgn. 52-24, Bldg. 201 (1 copy)
Mathematical Sciences Northwest, Inc., P.O. Box 1887, Bellevue, WA 98009 - Attn: Dr. Abraham Hertzberg (1 copy)
Martin Marietta Corp., P.O. Box 179, Mail Station 0471, Denver, CO 80201 - Attn: Mr. Stewart Chapin (1 copy)
Massachusetts Institute of Technology, Lincoln Laboratory, P.O. Box 73, Lexington, MA 02173 - Attn: Dr. S. Edelberg (1 copy)
Dr. L. C. Marquet (1 copy)
McDonnell Douglas Astronautics Co., 5301 Bolsa Avenue, Huntington Beach, CA 92647 - Attn: Mr. P. L. Klevatt, Dept. A3-830-BBFO, M/S 9 (1 copy)
McDonnell Douglas Research Labs., Dept. 220, Box 516, St. Louis, MO 63166 - Attn: Dr. D. P. Ames (1 copy)
MITRE Corp., P.O. Box 208, Bedford, MA 01730 - Attn: Mr. A. C. Cron (1 copy)
North American Rockwell Corp., Autonetics Div., Anaheim, CA 92803 - Attn: Mr. T. T. Kumagi, C/476 Mail Code HA18 (1 copy)
Northrop Corp., 3401 West Broadway, Hawthorne, CA 90250 - Attn: Dr. Gerard Hasserjian, Laser Systems Dept. (1 copy)
Dr. Anthony N. Pirri, Physical Sciences, Inc., 18 Lakeside Office Park, Wakefield, MA 01880 (1 copy)
RAND Corp., 1700 Main Street, Santa Monica, CA 90406 - Attn: Dr. C. R. Culp/Mr. G. A. Carter (1 copy)
Raytheon Co., 28 Seyon Street, Waltham, MA 02154 - Attn: Dr. F. A. Horrigan (Res. Div.) (1 copy)
Raytheon Co., Boston Post Road, Sudbury, MA 01776 - Attn: Dr. C. Sonnenschien (Equip. Div.) (1 copy)
Raytheon Co., Bedford Labs, Missile Systems Div., Bedford, MA 01730 - Attn: Dr. H. A. Mehlhorn (1 copy)
Riverside Research Institute, 80 West End Street, New York, NY 10023 - Attn: Dr. L. H. O'Neill (1 copy)
Dr. John Bose (1 copy)
(HPEGL Library) (1 copy)
R&D Associates, Inc., P.O. Box 3580, Santa Monica, CA 90431 - Attn: Dr. R. E. LeLevier (1 copy)
Rockwell International Corporation, Rocketdyne Division, Albuquerque District Office, 3636 Menaul Blvd., NE, Suite 211, Albuquerque, NM 87110 - Attn: C. K. Kraus, Mgr. (1 copy)
SANDIA Corp., P.O. Box 5800, Albuquerque, NM 87115 - Attn: Dr. Al Narath (1 copy)
Stanford Research Institute, Menlo Park, CA 94025 - Attn: Dr. F. T. Smith (1 copy)
Science Applications, Inc., 1911 N. Ft. Meyer Drive, Arlington, VA 22209 - Attn: L. Peckam (1 copy)
Science Applications, Inc., P.O. Box 328, Ann Arbor, MI 48103 - Attn: R. E. Meredith (1 copy)
Science Applications, Inc., 6 Preston Court, Bedford, MA 01703 - Attn: R. Greenberg (1 copy)
Science Applications, Inc., P.O. Box 2351, La Jolla, CA 92037 - Attn: Dr. John Asmus (1 copy)
Systems, Science and Software, P.O. Box 1620, La Jolla, CA 92037 - Attn: Alan F. Klein (1 copy)
Systems Consultants, Inc., 1050 31st Street, NW, Washington, D.C. 20007 - Attn: Dr. R. B. Keller (1 copy)
Thiokol Chemical Corp., WASATCH Division, P.O. Box 524, Brigham City, UT 84302 - Attn: Mr. J. E. Hansen (1 copy)
TRW Systems Group, One Space Park, Bldg. R-1, Rm. 1050, Redondo Beach, CA 90278 - Attn: Mr. Norman Campbell (1 copy)
United Technologies Research Center, 400 Main Street, East Hartford, CT 06108 - Attn: Mr. G. H. McLafferty (3 copies)

DISTRIBUTION LIST (Continued)

United Technologies Research Center, Pratt and Whitney Aircraft Div., Florida R&D Center, West Palm Beach, FL 33402 Attn: Dr. R. A. Schmidtke (1 copy)
Mr. Ed Pinsley (1 copy)

VARIAN Associates, EIMAC Division, 301 Industrial Way, San Carlos, CA 94070 - Attn: Mr. Jack Quinn (1 copy)

Vought Systems Division, LTV Aerospace Corp., P.O. Box 5907, Dallas, TX 75222 - Attn: Mr. F.G. Simpson, MS 254142 (1 copy)

Westinghouse Electric Corp., Defense and Space Center, Balt-Wash. International Airport - Box 746, Baltimore, MD 21203 - Attn: Mr. W.F. List (1 copy)

Westinghouse Research Labs., Beulah Road, Churchill Boro, Pittsburgh, PA 15235 - Attn: Dr. E. P. Riedel (1 copy)

United Technologies Research Center, East Hartford, CT 06108 - Attn: A. J. DeMaria (1 copy)

Airborne Instruments Laboratory, Walt Whitman Road, Melville, NY 11746 - Attn: F. Pace (1 copy)

General Electric R&D Center, Schenectady, NY 12305 - Attn: Dr. Donald White (1 copy)

Cleveland State University, Cleveland, OH 44115 - Attn: Dean Jack Soules (1 copy)

EXXON Research and Engineering Co., P.O. Box 8, Linden, NJ 07036 - Attn: D. Grafstein (1 copy)

University of Maryland, Department of Physics and Astronomy, College Park, MD 20742 - Attn: D. Currie (1 copy)

Sylvania Electric Products, Inc., 100 Fergeson Drive, Mountain View, CA 94040 - Attn: L. M. Osterink (1 copy)

North American Rockwell Corp., Autonetics Division, 3370 Miraloma Avenue, Anaheim, CA 92803 - Attn: R. Gudmundsen (1 copy)

Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02138 - Attn: Prof. A. Javan (1 copy)

Lockheed Missile & Space Co., Palo Alto Research Laboratories, Palo Alto, CA 94304 - Attn: Dr. R. C. Ohlman (1 copy)

ILC Laboratories, Inc., 164 Commercial Street, Sunnyvale, CA 94086 - Attn: L. Noble (1 copy)

University of Texas at Dallas, P.O. Box 30365, Dallas, TX 75230 - Attn: Prof. Carl B. Collins (1 copy)

Polytechnic Institute of New York, Rt. 110, Farmingdale, NY 11735 - Attn: Dr. William T. Walter (1 copy)

BEST AVAILABLE COPY